



On zeros of analytic functions satisfying non-radial growth conditions

Alexander Borichev, Leonid Golinskii and Stanislav Kupin

Abstract. Extending the results of Borichev–Golinskii–Kupin (2009), we obtain refined Blaschke-type necessary conditions on the zero distribution of analytic functions on the unit disk and on the complex plane with a cut along the positive semi-axis satisfying some non-radial growth restrictions.

To Peter Yuditskii on occasion of his 60-th anniversary

1. Introduction and main results

We have numbered
this first section

The study of relations between the zero distribution of an analytic function and its growth is likely to be one of the most basic problems of complex analysis. We have no intention to review a vast literature on it, but just give several references related to the points of our interest. Perhaps, the first results in this direction were obtained in the second half of 19th century by Hadamard, Borel, Weierstrass and others, see Levin Chapter 1 of [22] for a modern presentation. These results completely described the behavior of zeros of an entire function of finite type. Later, Blaschke [2], Nevanlinna [23] and Smirnov [27] described the zero sets of functions from the Hardy spaces $H^p(\mathbb{D})$, $p > 0$, or, more generally, the Nevanlinna class $\mathcal{N}(\mathbb{D})$. Here, as usual, $\mathbb{D} = \{|z| < 1\}$. Namely, for $f \in \mathcal{N}(\mathbb{D})$, $f \not\equiv 0$, one has

$$(1.1) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|) \leq \sup_{0 \leq r < 1} \frac{1}{2\pi} \int_0^{2\pi} \log^+ |f(re^{i\theta})| d\theta - \log |f(0)|,$$

where $Z(f)$ stands for the zero set of f counting multiplicities. Hence, a discrete subset $Z(f)$ of the unit disk is a zero set of a function from $H^p(\mathbb{D})$ (or $\mathcal{N}(\mathbb{D})$) if and only if the series at the LHS of (1.1) converges. This condition is usually called the “Blaschke condition” after [2].

Mathematics Subject Classification (2010): Primary 30C15; Secondary 30C20.

Keywords: Zeros of analytic functions, generalized Blaschke condition, Jensen-type formulae.

Let $\mathcal{A}(\mathbb{D})$ be the set of analytic functions on the unit disk. An argument similar to the proof of (1.1) shows that if $f \in \mathcal{A}(\mathbb{D})$, $|f(0)| = 1$, satisfies the growth condition

$$\log |f(z)| \leq \frac{K}{(1 - |z|)^p},$$

where $p \geq 1$, then for any $\varepsilon > 0$,

$$(1.2) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|)^{p+1+\varepsilon} \leq C_0 \cdot K,$$

where the constant $C_0 = C_0(p, \varepsilon)$ depends on p and ε , see, e.g., Golubev [16].

Of course, the study of the zero distribution of analytic functions from other classes is much more involved; see, for instance, papers of Korenblum [18], [19] on the zero distribution for functions from spaces $A^{-p}(\mathbb{D})$, $A^{-\infty}(\mathbb{D})$. Interesting results on zeros of functions from some Bergman-type spaces are given in Seip [26].

The above mentioned spaces of analytic functions are defined with the help of a radial (i.e., invariant with respect to rotations of the unit disk) growth conditions. However, it turns out that one often needs to deal with classes of analytic functions subject to *non-radial* growth relations. These classes appear, in particular, if one wants to study the distribution of the discrete spectrum of non-self-adjoint perturbations for certain self-adjoint or unitary operators.

The study of such classes was initiated in [3], and the main result therein looks as follows, see [3], Theorem 0.2. Given a finite set $F = \{\xi_k\}_{k=1}^m$ on the unit circle $\mathbb{T} = \{|z| = 1\}$, let $d(z, F) = \min_k |z - \xi_k|$ denote the Euclidian distance between a point $z \in \mathbb{D}$ and F . In what follows, $a_+ := \max(a, 0)$, $a_- := \max(-a, 0)$, and K is a positive constant.

Theorem A. *Let $f \in \mathcal{A}(\mathbb{D})$, $|f(0)| = 1$, satisfy the growth condition*

$$\log |f(z)| \leq \frac{K}{(1 - |z|)^p d^q(z, F)}, \quad z \in \mathbb{D}, \quad p, q \geq 0.$$

Then, for each $\varepsilon > 0$ there is a positive number $C_1 = C_1(F, p, q, \varepsilon)$ such that the following Blaschke-type condition holds:

$$(1.3) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|)^{p+1+\varepsilon} d^{(q-1+\varepsilon)_+}(\zeta, F) \leq C_1 \cdot K.$$

Moreover, in the case $p = 0$ the term $(1 - |\zeta|)^{p+1+\varepsilon}$ can be replaced by $(1 - |\zeta|)$.

Theorem A effectively applies to the study of the discrete spectrum of complex perturbations of certain self-adjoint operators of mathematical physics in Demuth–Hansmann–Katriel [5], [6], Golinskii–Kupin [13], [14], [15], Dubuisson [7], [8], and Sambou [25]. We also mention recent interesting papers by Cuenin–Laptev–Tretter [4], Frank–Sabin [11], Frank [12], and Laptev–Safronov [20] in this connection. For some extensions of this result to the case of arbitrary closed sets F and subharmonic on \mathbb{D} functions f , and applications in perturbation theory see Favorov–Golinskii [9], [10].

Let us go over to the main results of the present paper which extend Theorem A. Let $E = \{\zeta_j\}_{j=1}^n$ and $F = \{\xi_k\}_{k=1}^m$ be two disjoint finite sets of distinct points on the unit circle \mathbb{T} .

Theorem 1.1. *Let $f \in \mathcal{A}(\mathbb{D})$, $|f(0)| = 1$, satisfy the growth condition*

$$(1.4) \quad \log |f(z)| \leq \frac{K}{(1 - |z|)^p} \frac{d^r(z, E)}{d^q(z, F)}, \quad z \in \mathbb{D}, \quad p, q, r \geq 0.$$

Then for every $\varepsilon > 0$, there is a positive number $C_2 = C_2(E, F, p, q, r, \varepsilon)$ such that the following Blaschke-type condition holds:

$$(1.5) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|)^{p+1+\varepsilon} \frac{d^{(q-1+\varepsilon)_+}(\zeta, F)}{d^{\min(p,r)}(\zeta, E)} \leq C_2 \cdot K.$$

Of course, Theorem A is exactly Theorem 1.1 with $r = 0$.

An obvious inequality for an arbitrary finite set $B = \{\beta_j\}_{j=1}^n \subset \mathbb{T}$,

$$c(B) \prod_{j=1}^n |z - \beta_j| \leq d(z, B) \leq C(B) \prod_{j=1}^n |z - \beta_j|,$$

along with Theorem 0.3 from [3] prompt a more general statement.

Theorem 1.2. *Let $f \in \mathcal{A}(\mathbb{D})$, $|f(0)| = 1$, satisfy the growth condition*

$$(1.6) \quad \log |f(z)| \leq \frac{K}{(1 - |z|)^p} \frac{\prod_{j=1}^n |z - \zeta_j|^{r_j}}{\prod_{k=1}^m |z - \xi_k|^{q_k}}, \quad z \in \mathbb{D}, \quad p, q_k, r_j \geq 0.$$

Then for every $\varepsilon > 0$, there is a positive number $C_3 = C_3(E, F, p, \{q_k\}, \{r_j\}, \varepsilon)$ such that the following Blaschke-type condition holds:

$$(1.7) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|)^{p+1+\varepsilon} \frac{\prod_{k=1}^m |\zeta - \xi_k|^{(q_k-1+\varepsilon)_+}}{\prod_{j=1}^n |\zeta - \zeta_j|^{\min(p,r_j)}} \leq C_3 \cdot K.$$

Once again, in the case $p = 0$ the factor $(1 - |\zeta|)^{1+\varepsilon}$ in (1.5) and (1.7) can be replaced by $(1 - |\zeta|)$.

Remark. An observation due to Hansmann–Katriel [17] applies in our setting. It turns out that the stronger assumption

$$\log |f(z)| \leq \frac{K|z|^\gamma}{(1 - |z|)^p} \frac{\prod_{j=1}^n |z - \zeta_j|^{r_j}}{\prod_{k=1}^m |z - \xi_k|^{q_k}}, \quad z \in \mathbb{D}, \quad p, q_k, r_j, \gamma \geq 0,$$

implies the stronger conclusion

$$\sum_{\zeta \in Z(f)} \frac{(1 - |\zeta|)^{p+1+\varepsilon}}{|\zeta|^{(\gamma-\varepsilon)_+}} \frac{\prod_{k=1}^m |\zeta - \xi_k|^{(q_k-1+\varepsilon)_+}}{\prod_{j=1}^n |\zeta - \zeta_j|^{\min(p,r_j)}} \leq C(E, F, p, \{q_k\}, \{r_j\}, \varepsilon) \cdot K.$$

The result of Theorem 1.1 can be extended in another direction involving arbitrary closed subsets F of the unit circle. A key ingredient in such extensions is the following quantitative characteristic of F known as the *Ahern–Clark type* [1]:

$$\alpha(F) := \sup \{ \alpha \in \mathbb{R} : |\{t \in \mathbb{T} : d(t, F) < x\}| = O(x^\alpha), x \rightarrow +0 \}.$$

Here $|A|$ denotes the Lebesgue measure of a measurable set $A \subset \mathbb{T}$.

Theorem 1.3. *Let $E = \{\zeta_j\}_{j=1}^n$ be a finite subset of \mathbb{T} , $F \subset \mathbb{T}$ be an arbitrary closed set, and $E \cap F = \emptyset$. Let $f \in \mathcal{A}(\mathbb{D})$, $|f(0)| = 1$, satisfy the growth condition (1.4). Then for every $\varepsilon > 0$, there is a positive number $C_4 = C_4(E, F, p, q, r, \varepsilon)$ such that the following Blaschke-type condition holds:*

$$(1.8) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|)^{p+1+\varepsilon} \frac{d^{(q-\alpha(F)+\varepsilon)_+}(\zeta, F)}{d^{\min(p,r)}(\zeta, E)} \leq C_4 \cdot K.$$

Clearly, Theorem 1.1 is a special case of the latter result, since $\alpha(F) = 1$ for finite sets F .

As we will see later in Section 4, inequalities (1.5), (1.7) are in some sense “local” with respect to the singular points $\{\zeta_j\}_{j=1}^n$ and $\{\xi_k\}_{k=1}^m$ on the unit circle, so we can restrict ourselves to the case $n = m = 1$ and $E = \{\zeta_0\}$, $F = \{\xi_0\}$. The following “one-point” version of the main result will be crucial in the sequel.

Theorem 1.4. *Let $\zeta_0, \xi_0 \in \mathbb{T}$, $\zeta_0 \neq \xi_0$, and let $f \in \mathcal{A}(\mathbb{D})$, $|f(0)| = 1$, satisfy the growth condition*

$$(1.9) \quad \log |f(z)| \leq \frac{K}{(1 - |z|)^p} \frac{|z - \zeta_0|^r}{|z - \xi_0|^q}, \quad z \in \mathbb{D}, \quad p, q, r \geq 0.$$

Then for every $\varepsilon > 0$, there is a positive number $C_5 = C_5(\zeta_0, \xi_0, p, q, r, \varepsilon)$ such that the following inequality holds:

$$(1.10) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|)^{p+1+\varepsilon} \frac{|\zeta - \xi_0|^{(q-1+\varepsilon)_+}}{|\zeta - \zeta_0|^{\min(p,r)}} \leq C_5 \cdot K.$$

The paper is organized in a straightforward manner. The preliminaries are given in Section 2. In Sections 3 and 4 we prove Theorem 1.4 and then deduce the general statements in Theorems 1.2 and 1.3 from this one-point version. Some further results (the analogs for the upper half-plane and the plane with a cut) are given in Section 5.

To keep the notation reasonably simple and consistent, we usually number the constants C_k appearing in the formulations of theorems, propositions, etc. The constants C arising in the proofs are generic, i.e., the same symbol does not necessarily denote the same constant in different occurrences.

2. Conformal mappings, Pommerenke lemma and Stolz angles

We start with some general preliminaries from complex analysis.

The known distortion inequalities [24], Corollary 1.4, play a key role in what follows.

Lemma 2.1. *Let Ω be a bounded, simply connected domain with the boundary $\partial\Omega$, and φ be a conformal mapping of Ω onto \mathbb{D} . Then*

$$(2.1) \quad \frac{1}{2} d(w, \partial\Omega) \cdot |\varphi'(w)| \leq 1 - |\varphi(w)| \leq 4 d(w, \partial\Omega) \cdot |\varphi'(w)|, \quad w \in \Omega.$$

This result will be applied in the following situation, wherein the bounds on derivatives can be specified. It is related to the Stolz angle with the vertex at $\zeta_0 \in \mathbb{T}$, that is, a domain inside the unit disk of the form

$$(2.2) \quad \mathcal{S}_A(\zeta_0) := \left\{ z \in \mathbb{D} : \frac{|z - \zeta_0|}{1 - |z|} < A \right\}, \quad A > 1.$$

When $\zeta_0 = 1$, we use the abbreviation $\mathcal{S}_A := \mathcal{S}_A(1)$, see Figure 1. The interior angle of \mathcal{S}_A at 1 equals 2ω , $0 < \omega := \arccos A^{-1} < \pi/2$. The Stolz angles $\{\mathcal{S}_A\}_{A>1}$ form an increasing family of sets which exhaust the unit disk as $A \rightarrow \infty$. The boundary of \mathcal{S}_A is




FIGURE 1. Stolz angle $\mathcal{S}_A = \mathcal{S}_A(1)$, $A > 1$, $\omega = \arccos A^{-1}$.

Let φ_A denote the conformal mapping $\varphi_A : \mathcal{S}_A \rightarrow \mathbb{D}$, $\varphi_A(0) = 0$, $\varphi_A(1) = 1$. The following result provides a local uniform bound for its derivative φ'_A .

Lemma 2.2. *Let*

$$\alpha = \alpha_A := \frac{\pi}{2\omega} = \frac{\pi}{2 \arccos A^{-1}}, \quad \alpha > 1.$$

Then the following bounds hold uniformly for $A \geq 2$:

$$(2.3) \quad \frac{1}{16} < \frac{|\varphi'_A(z)|}{|z - 1|^{\alpha-1}} < 48, \quad z \in \mathcal{S}_A^+ := \mathcal{S}_A \cap \left\{ |z - 1| < \frac{1}{16} \right\}.$$

Proof. We just sketch the proof, which is rather standard. Let $\psi: \mathbb{D} \rightarrow \mathbb{C}_r := \{z : \operatorname{Re} z > 0\}$ be the linear-fractional mapping of \mathbb{D} onto the right half-plane, $\psi(0) = 1$, $\psi(1) = \infty$. A crucial observation is that ψ maps \mathcal{S}_A onto the interior H_i of the right branch of the hyperbola¹

$$H : \frac{x^2}{\cos^2 \omega} - \frac{y^2}{\sin^2 \omega} = 1, \quad z = x + iy.$$

Set $\mathbb{C}_\pm = \{z : \pm \operatorname{Im} z > 0\}$, $H_\pm = H_i \cap \mathbb{C}_\pm$, and define $\phi_1(z) = z + \sqrt{z^2 - 1} = \exp(\operatorname{arch} z)$, $\phi_1: H_\pm \rightarrow A_\pm = \{re^{i\theta} : r > 1, \pm\theta \in (0, \omega)\}$, $\phi_2(z) = (z^{\pi/\omega} + z^{-\pi/\omega})/2$, $\phi_2: A_\pm \rightarrow \mathbb{C}_\pm$, $\phi_3(z) = \sqrt{1+z}$, $\phi_3: \mathbb{C}_\pm \rightarrow \mathbb{C}_\pm \cap \mathbb{C}_r$. Since $\phi_3 \circ \phi_2 \circ \phi_1$ extends continuously to $H_i \cap \mathbb{R}$, we obtain a conformal map $\phi: H_i \rightarrow \mathbb{C}_r$, $\phi(1) = \sqrt{2}$, $\phi(\infty) = \infty$. Finally, if $\phi_4(z) = (z - \sqrt{2})/(z + \sqrt{2})$, then

$$\varphi_A = \phi_4 \circ \phi \circ \psi = \left(\frac{(1 \pm \sqrt{z})^\alpha - (1 \mp \sqrt{z})^\alpha}{(1 \pm \sqrt{z})^\alpha + (1 \mp \sqrt{z})^\alpha} \right)^2 = 1 - \frac{4(1-z)^\alpha}{((1 \pm \sqrt{z})^\alpha + (1 \mp \sqrt{z})^\alpha)^2}$$

(see also Lavrent'ev–Shabat Chapter 2.3.36 in [21]). Next,

$$(2.4) \quad |\varphi'_A(z)| = \frac{4\alpha|1-z|^{\alpha-1}}{\sqrt{|z|}} \cdot \frac{|(1 \pm \sqrt{z})^\alpha - (1 \mp \sqrt{z})^\alpha|}{|(1 \pm \sqrt{z})^\alpha + (1 \mp \sqrt{z})^\alpha|^3}.$$

Since $\alpha \leq 3/2$ for $A \geq 2$, the elementary bounds

$$\begin{aligned} \frac{1}{2} &\leq \sqrt{|z|} \leq 1, \\ 1 &\leq |(1 \pm \sqrt{z})^\alpha + (1 \mp \sqrt{z})^\alpha| \leq 4, \\ 1 &\leq |(1 \pm \sqrt{z})^\alpha - (1 \mp \sqrt{z})^\alpha| \leq 4, \end{aligned}$$

valid for $z \in \mathcal{S}_A^+$, yield (2.3). □

The following simple relation between two Stolz angles is casted as a lemma for convenience only; its elementary proof is omitted.

Lemma 2.3. *Let $A < B$, so $\mathcal{S}_A \subset \mathcal{S}_B$. Then, for $z \in \mathcal{S}_A$,*

$$(2.5) \quad \frac{B-A}{B+1} (1-|z|) \leq d(z, \partial\mathcal{S}_B) < 1-|z|.$$

For $0 < a < 1$, consider a nested family of domains (curvilinear quadrangles) $\{L_a\}$, see Figure 2,

$$(2.6) \quad L_{a_1} \subset L_{a_2} \subset \mathbb{D}, \quad 0 < a_1 < a_2 < 1.$$

We denote by $\eta = \eta_a$ the conformal mapping of L_a onto \mathbb{D} with normalization $\eta(0) = 0$, $\eta(1) = 1$, and write η_j , $j = 1, 2$ for the domains L_{a_j} . Although there

¹We thank D. Tulyakov for this remark.

is no explicit formula for η , it is easily seen from Theorem 3.9 in [24] that both η and η' have continuous extensions on the closure $\overline{L_a}$, and so

$$(2.7) \quad |\eta'(z)| \leq c(a), \quad z \in L_a.$$

The relations below follow directly from (2.7) and Lemma 2.1. First,

$$(2.8) \quad 1 - |\eta_2(z)| \leq c_1(a_2)(1 - |z|), \quad z \in L_{a_2},$$

holds with some positive constant $c_1(a_2)$. Next, since 1 is a regular point for η (η is analytic at some neighborhood of 1),

$$(2.9) \quad c_2(a_2) \leq \frac{|1 - \eta_2(z)|}{|1 - z|} \leq c_3(a_2), \quad z \in L_{a_2},$$

holds with some positive constants $c_j(a_2)$, $j = 2, 3$. Finally, there are positive constants $c_j = c_j(a_1, a_2)$, $j = 4, 5$, such that

$$(2.10) \quad c_4(a_1, a_2) \leq \frac{1 - |\eta_2(z)|}{1 - |z|} \leq c_5(a_1, a_2), \quad z \in L_{a_1}.$$

We will exploit these relations later in Section 4.




FIGURE 2. Domains $L_{a_1}, L_{a_2}, 0 < a_1 < a_2$.

3. Proof of Theorem 1.4 for $q = 0$

Without loss of generality we assume that $\zeta_0 = 1$. By (1.9),

$$\log |f(z)| \leq \frac{2^r K}{(1 - |z|)^p}, \quad z \in \mathbb{D},$$

and, by (1.2), for each $\varepsilon > 0$,

$$(3.1) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|)^{p+1+\varepsilon} \leq C(p, r, \varepsilon) \cdot K.$$

To clarify the local character of the problem, put

$$(3.2) \quad Z^+(f) := Z(f) \cap \left\{ |z - 1| < \frac{1}{16} \right\}, \quad Z^-(f) := Z(f) \cap \left\{ |z - 1| \geq \frac{1}{16} \right\},$$

so that with $s := \min(p, r)$ we have

$$\sum_{\zeta \in Z(f)} \frac{(1 - |\zeta|)^{p+1+\varepsilon}}{|1 - \zeta|^s} = \left\{ \sum_{\zeta \in Z^+(f)} + \sum_{\zeta \in Z^-(f)} \right\} \frac{(1 - |\zeta|)^{p+1+\varepsilon}}{|1 - \zeta|^s} = \Sigma^+ + \Sigma^-.$$

The bound for Σ^- follows directly from (3.1) and the inequality $|1 - \zeta| \geq 1/16$:

$$(3.3) \quad \Sigma^- = \sum_{\zeta \in Z^-(f)} \frac{(1 - |\zeta|)^{p+1+\varepsilon}}{|1 - \zeta|^s} \leq C(p, r, \varepsilon) \cdot K, \quad p, r \geq 0.$$

Thus, the main problem is to prove (1.10) for Σ^+ .

3.1. Case $p \leq r$

Given a function $f \in \mathcal{A}(\mathbb{D})$ and a number $A \geq 2$, put (see (3.2) and (2.3))

$$(3.4) \quad Z_A(f) := Z^+(f) \cap \mathcal{S}_A = Z(f) \cap \mathcal{S}_A^+.$$

Step 1. Let $\psi_A = \varphi_A^{(-1)}$ be the conformal mapping from \mathbb{D} onto \mathcal{S}_A , $\psi_A(0) = 0$. Set $f_A = f(\psi_A)$. Then $f_A \in \mathcal{A}(\mathbb{D})$, $|f_A(0)| = 1$ and by (1.9) we have

$$\log |f_A(w)| \leq K \frac{|1 - \psi_A(w)|^r}{(1 - |\psi_A(w)|)^p} \leq 2^{r-p} A^p \cdot K, \quad w \in \mathbb{D}.$$

The Poisson–Jensen formula implies

$$(3.5) \quad \sum_{w \in Z(f_A)} (1 - |w|) \leq 2^{r-p} A^p \cdot K.$$

However, $Z(f_A) = \varphi_A(Z(f) \cap \mathcal{S}_A)$, and so

$$(3.6) \quad \sum_{w \in Z(f_A)} (1 - |w|) = \sum_{\zeta \in Z(f) \cap \mathcal{S}_A} (1 - |\varphi_A(\zeta)|) \leq 2^{r-p} A^p \cdot K.$$

By Lemma 2.1,

$$\sum_{\zeta \in Z(f) \cap \mathcal{S}_A} |\varphi'_A(\zeta)| \cdot d(\zeta, \partial \mathcal{S}_A) \leq 2^{r-p+1} A^p \cdot K,$$

and, since $Z(f) \cap \mathcal{S}_A \supset Z_A(f) = Z(f) \cap \mathcal{S}_A^+$, it follows from Lemma 2.2 that for $A \geq 2$

$$\begin{aligned} \sum_{\zeta \in Z(f) \cap \mathcal{S}_A} |\varphi'_A(\zeta)| \cdot d(\zeta, \partial \mathcal{S}_A) &\geq \sum_{\zeta \in Z_A(f)} |\varphi'_A(\zeta)| \cdot d(\zeta, \partial \mathcal{S}_A) \\ &\geq \frac{1}{32} \sum_{\zeta \in Z_A(f)} |\zeta - 1|^{\alpha-1} \cdot d(\zeta, \partial \mathcal{S}_A). \end{aligned}$$

Hence,

$$(3.7) \quad \sum_{\zeta \in Z_A(f)} |1 - \zeta|^{\alpha-1} \cdot d(\zeta, \partial S_A) \leq 2^{r-p+6} A^p \cdot K, \quad \alpha = \frac{\pi}{2 \arccos A^{-1}}.$$

Step 2. In what follows $A = A_k = 2^k$, $k \in \mathbb{N}$, so the Stolz angles $S_k := S_{A_k}$ (with a little abuse of notation) exhaust the unit disk, as $k \rightarrow \infty$. Relation (3.7) with $A = A_{k+1}$ takes the form

$$\sum_{\zeta \in Z_{k+1}} |1 - \zeta|^{\alpha_{k+1}-1} \cdot d(\zeta, \partial S_{k+1}) \leq 2^{kp+r+6} K, \quad Z_k := Z_{A_k}(f) = Z^+(f) \cap S_k,$$

see (3.2), (3.4), or, since $Z_k \subset Z_{k+1}$,

$$(3.8) \quad \sum_{\zeta \in Z_k} |1 - \zeta|^{\beta_{k+1}} \cdot d(\zeta, \partial S_{k+1}) \leq 2^{kp+r+6} K, \quad \beta_{k+1} := \alpha_{k+1} - 1.$$

To apply Lemma 2.3 with $A = 2^k$, $B = 2^{k+1}$, notice that

$$\frac{B - A}{B + 1} = \frac{2^{k+1} - 2^k}{2^{k+1} + 1} \geq \frac{2}{5},$$

so (3.8) entails

$$(3.9) \quad \sum_{\zeta \in Z_k} (1 - |\zeta|) |1 - \zeta|^{\beta_{k+1}} \leq 5K \cdot 2^{kp+r+5} = C(r) 2^{kp} \cdot K,$$

for $k \in \mathbb{N}$. It is convenient to deal with a chain of inequalities

$$\sum_{\zeta \in Z_k \setminus Z_{k-1}} (1 - |\zeta|) |1 - \zeta|^{\beta_{k+1}} \leq C(r) 2^{kp} \cdot K, \quad k \in \mathbb{N}, \quad Z_0 := \emptyset.$$

Take an arbitrary $0 < \varepsilon < 1/16$ and write

$$(3.10) \quad \frac{1}{2^{k(p+\varepsilon)}} \sum_{\zeta \in Z_k \setminus Z_{k-1}} (1 - |\zeta|) |1 - \zeta|^{\beta_{k+1}} \leq C(r) 2^{-\varepsilon k} \cdot K.$$

On the set $Z_k \setminus Z_{k-1}$ we have

$$2^{-k} < \frac{1 - |\zeta|}{|1 - \zeta|} \leq 2^{-k+1}, \quad \left(\frac{1 - |\zeta|}{|1 - \zeta|} \right)^{p+\varepsilon} \leq \frac{2^{p+\varepsilon}}{2^{k(p+\varepsilon)}},$$

and so

$$(3.11) \quad \sum_{\zeta \in Z_k \setminus Z_{k-1}} \frac{(1 - |\zeta|)^{p+\varepsilon+1}}{|1 - \zeta|^{p+\varepsilon}} |1 - \zeta|^{\beta_{k+1}} \leq C(p, r) 2^{-\varepsilon k} \cdot K.$$

Step 3. We have

$$\alpha_k = \frac{\pi}{2 \arccos 2^{-k}}, \quad \beta_k = \alpha_k - 1 = \frac{\arcsin 2^{-k}}{\arccos 2^{-k}},$$

and as $x \leq \arcsin x \leq \pi x/2$ for $0 \leq x \leq 1$, and $\arccos 1/2 = \pi/3$, we see that

$$(3.12) \quad \frac{2}{\pi} \leq 2^k \beta_k \leq \frac{3}{2}.$$

By definition, $\beta_k \searrow 0$ as $k \rightarrow \infty$. Now, choose $k_0 = k_0(\varepsilon)$ from the relations

$$(3.13) \quad 2^{-k_0-1} \leq \varepsilon < 2^{-k_0},$$

and hence

$$(3.14) \quad \mathcal{S}_{k_0} \subset \mathcal{S}_{1/\varepsilon} \subset \mathcal{S}_{k_0+1}, \quad Z_{k_0} \subset Z^+(f) \cap \mathcal{S}_{1/\varepsilon} \subset Z_{k_0+1}.$$

By (3.12) and (3.13), one has for $k \geq k_0 + 1$

$$\beta_{k+1} \leq \beta_{k_0+2} \leq \frac{3}{2} 2^{-k_0-2} < \frac{3}{4} \varepsilon.$$

Let $z \in Z^+(f)$. Since $|1 - z| < 1/16$, we see that $|1 - z|^{\beta_{k+1}} \geq |1 - z|^\varepsilon$. Hence, (3.11) implies that

$$(3.15) \quad \sum_{\zeta \in Z_k \setminus Z_{k-1}} \frac{(1 - |\zeta|)^{p+\varepsilon+1}}{|1 - \zeta|^p} \leq C(p, r) 2^{-\varepsilon k} \cdot K, \quad k \geq k_0 + 1.$$

Summation over k from $k = k_0 + 1$ to infinity gives

$$(3.16) \quad \sum_{\zeta \in Z^+(f) \setminus Z_{k_0}} \frac{(1 - |\zeta|)^{p+\varepsilon+1}}{|1 - \zeta|^p} \leq C(p, r, \varepsilon) \cdot K.$$

Next, write

$$Z^+(f) = (Z^+(f) \cap \mathcal{S}_{1/\varepsilon}) \cup (Z^+(f) \cap \mathcal{S}_{1/\varepsilon}^c), \quad \mathcal{S}_{1/\varepsilon}^c := \mathbb{D} \setminus \mathcal{S}_{1/\varepsilon}.$$

By (3.14), $Z^+(f) \setminus Z_{k_0} \supset Z^+(f) \cap \mathcal{S}_{1/\varepsilon}^c$, so (3.16) provides

$$(3.17) \quad \sum_{\zeta \in Z^+(f) \cap \mathcal{S}_{1/\varepsilon}^c} \frac{(1 - |\zeta|)^{p+\varepsilon+1}}{|1 - \zeta|^p} \leq C(p, r, \varepsilon) \cdot K.$$

On the other hand, put $k = k_0 + 1$ in (3.9). By (3.13), $\beta_{k_0+2} < \varepsilon$, and (3.14) implies that

$$(3.18) \quad \sum_{\zeta \in Z^+(f) \cap \mathcal{S}_{1/\varepsilon}} (1 - |\zeta|)|1 - \zeta|^\varepsilon \leq \sum_{\zeta \in Z_{k_0+1}} (1 - |\zeta|)|1 - \zeta|^\varepsilon \leq C(r, \varepsilon) \cdot K.$$

The sum of (3.17) and (3.18) gives

$$(3.19) \quad \sum_{\zeta \in Z^+(f) \cap \mathcal{S}_{1/\varepsilon}^c} \frac{(1 - |\zeta|)^{p+\varepsilon+1}}{|1 - \zeta|^p} + \sum_{\zeta \in Z^+(f) \cap \mathcal{S}_{1/\varepsilon}} (1 - |\zeta|)|1 - \zeta|^\varepsilon \leq C(p, r, \varepsilon) \cdot K,$$

and it remains to note again that the inequality $|1 - z| \geq 1 - |z|$ for $z \in \mathbb{D}$ implies

$$(1 - |z|)|1 - z|^\varepsilon \geq \frac{(1 - |z|)^{p+\varepsilon+1}}{|1 - z|^p}.$$

Finally,

$$(3.20) \quad \Sigma^+ = \sum_{\zeta \in Z^+(f)} \frac{(1 - |\zeta|)^{p+\varepsilon+1}}{|1 - \zeta|^p} \leq C(p, r, \varepsilon) \cdot K.$$

Note that now $p = s = \min(p, r)$. A combination of (3.20) and (3.3) completes the proof of Theorem 1.4 in the case $q = 0, p \leq r$.

3.2. Case $p > r$

Let $f \in \mathcal{A}(\mathbb{D}), |f(0)| = 1$, satisfy

$$(3.21) \quad \log |f(z)| \leq K \frac{|1 - z|^r}{(1 - |z|)^p}, \quad z \in \mathbb{D},$$

with $0 \leq r < p$. Recalling the notation $f_A = f(\psi_A)$ (see Section 3.1) we have

$$\log |f_A(w)| \leq K \frac{|1 - \psi_A(w)|^r}{(1 - |\psi_A(w)|)^r} \cdot \frac{1}{(1 - |\psi_A(w)|)^{p-r}} \leq \frac{K A^r}{(1 - |\psi_A(w)|)^{p-r}}.$$

By the Schwarz lemma, $|\psi_A(w)| \leq |w|$, and so

$$(3.22) \quad \log |f_A(w)| \leq \frac{K A^r}{(1 - |w|)^{p-r}}.$$

As above in (3.1), we get for each $\varepsilon > 0$

$$(3.23) \quad \sum_{w \in Z(f_A)} (1 - |w|)^\gamma \leq C(p, r, \varepsilon) A^r \cdot K,$$

where $\gamma = \gamma(p, r, \varepsilon) := p - r + 1 + \varepsilon$. So we come to (3.5) with exponent γ instead of 1.

The rest is essentially the same as in the argument for the case $p \leq r$. For instance, (3.7) becomes

$$(3.24) \quad \sum_{\zeta \in Z_A(f)} |1 - \zeta|^{\gamma(\alpha-1)} \cdot d^\gamma(\zeta, \partial \mathcal{S}_A) \leq C(p, r) A^r \cdot K,$$

and (3.11) turns into

$$(3.25) \quad \sum_{\zeta \in Z_k \setminus Z_{k-1}} \frac{(1 - |\zeta|)^{p+1+2\varepsilon}}{|1 - \zeta|^{r+\varepsilon}} |1 - \zeta|^{\gamma\beta_{k+1}} \leq C(p, r) 2^{-\varepsilon k} \cdot K.$$

The choice of k_0 is somewhat different from (3.13):

$$2^{-k_0-1} \leq \frac{\varepsilon}{p - r + 2} < 2^{-k_0},$$

and again $\gamma\beta_{k+1} \leq \varepsilon$ for $k \geq k_0 + 1$. Thereby we come to

$$(3.26) \quad \sum_{\zeta \in Z^+(f) \setminus Z_{k_0}} \frac{(1 - |\zeta|)^{p+1+2\varepsilon}}{|1 - \zeta|^r} \leq C(p, r, \varepsilon) \cdot K;$$

compare this inequality to (3.16). Finally,

$$(3.27) \quad \sum_{\zeta \in Z^+(f)} \frac{(1 - |\zeta|)^{p+1+2\varepsilon}}{|1 - \zeta|^r} \leq C(p, r, \varepsilon) \cdot K.$$

A combination of (3.27) and (3.3) completes the proof of Theorem 1.4 for $q = 0$.

4. Proofs of Theorem 1.4 with $q > 0$, Theorems 1.2 and 1.3

We proceed with a local version of the result obtained in Section 3, see also Favorov–Golinskii [10].

Proposition 4.1. *Given the quadrangle L_{a_2} on Figure 2, let $g \in \mathcal{A}(L_{a_2})$ satisfy*

$$(4.1) \quad \log |g(w)| \leq K \frac{|1 - w|^r}{(1 - |w|)^p}, \quad w \in L_{a_2}, \quad p, r \geq 0.$$

Then for every $\varepsilon > 0$ and every $0 < a_1 < a_2$ there exists a positive constant $C = C(p, r, \varepsilon; a_1, a_2)$ such that

$$(4.2) \quad \sum_{\zeta \in Z(g) \cap L_{a_1}} \frac{(1 - |\zeta|)^{p+1+\varepsilon}}{|\zeta - 1|^s} \leq C \cdot K, \quad s = \min(p, r).$$

Proof. Recall that η_2 stands for the normalized conformal map from L_{a_2} onto \mathbb{D} . Put $f := g \circ \eta_2^{-1}$, so

$$\log |f(z)| \leq K \frac{|1 - \eta_2^{-1}(z)|^r}{(1 - |\eta_2^{-1}(z)|)^p}, \quad z \in \mathbb{D}.$$

In view of (2.8), (2.9) we have

$$\log |f(z)| \leq CK \frac{|1 - z|^r}{(1 - |z|)^p}, \quad z \in \mathbb{D}, \quad p, r \geq 0.$$

By the result obtained in Section 3, for every $\varepsilon > 0$

$$\sum_{v \in Z(f)} \frac{(1 - |v|)^{p+1+\varepsilon}}{|1 - v|^s} = \sum_{\zeta \in Z(g)} \frac{(1 - |\eta_2(\zeta)|)^{p+1+\varepsilon}}{|1 - \eta_2(\zeta)|^s} \leq C(p, r, \varepsilon; a_2) \cdot K, \quad s = \min(p, r),$$

and moreover, for $0 < a_1 < a_2$

$$\sum_{\zeta \in Z(g) \cap L_{a_1}} \frac{(1 - |\eta_2(\zeta)|)^{p+1+\varepsilon}}{|1 - \eta_2(\zeta)|^s} \leq C(p, r, \varepsilon; a_1, a_2) \cdot K.$$

The result now follows from (2.9) and (2.10). □

Proof of Theorem 1.4. Recall that, by convention, $\zeta_0 = 1$. To complete the proof of Theorem 1.4, we note that (1.9) implies (4.1) locally inside the domain L_a with $4a = |1 - \zeta_0|$ and with K replaced by $C(\zeta_0, \xi_0) \cdot K$. Put $\rho := (q - 1 + \varepsilon)_+$. By Proposition 4.1,

$$(4.3) \quad \sum_{\zeta \in Z(f) \cap L_{a/2}} (1 - |\zeta|)^{p+1+\varepsilon} \frac{|\zeta - \xi_0|^\rho}{|\zeta - 1|^s} \leq 2^\rho \sum_{\zeta \in Z(f) \cap L_{a/2}} \frac{(1 - |\zeta|)^{p+1+\varepsilon}}{|\zeta - 1|^s} \leq C(p, q, r, \xi_0, \varepsilon) \cdot K.$$

On the other hand, condition (1.9) implies the global bound

$$\log |f(z)| \leq \frac{2^r K}{(1 - |z|)^p |z - \xi_0|^q}, \quad z \in \mathbb{D},$$

and so

$$(4.4) \quad \begin{aligned} \sum_{\zeta \in Z(f) \setminus L_{a/2}} (1 - |\zeta|)^{p+1+\varepsilon} \frac{|\zeta - \xi_0|^\rho}{|\zeta - 1|^s} &\leq C \sum_{\zeta \in Z(f) \setminus L_{a/2}} (1 - |\zeta|)^{p+1+\varepsilon} |\zeta - \xi_0|^\rho \\ &\leq C \sum_{\zeta \in Z(f)} (1 - |\zeta|)^{p+1+\varepsilon} |\zeta - \xi_0|^\rho \leq C(p, q, r, \xi_0, \varepsilon) \cdot K. \end{aligned}$$

The latter inequality follows from Theorem A. The combination of (4.3) and (4.4) completes the proof of Theorem 1.4. \square

Proof of Theorem 1.2. We follow the line of reasoning of the above proof. In view of (1.6) one has the bound, which holds inside the turned quadrangle,

$$L_a(\zeta_i) = \zeta_i L_a, \quad a := \frac{1}{2} \min_{1 \leq j \leq n} d(\zeta_j, E \setminus \{\zeta_j\}).$$

Precisely,

$$(4.5) \quad \log |f(z)| \leq C K \frac{|z - \zeta_i|^{r_i}}{(1 - |z|)^p}, \quad z \in L_a(\zeta_i), \quad i = 1, 2, \dots, n.$$

By Proposition 4.1, for $i = 1, 2, \dots, n$ and $s_i = \min(p, r_i)$

$$(4.6) \quad \sum_{\zeta \in Z(f) \cap L_{a/2}(\zeta_i)} \frac{(1 - |\zeta|)^{p+1+\varepsilon}}{\prod_{j=1}^n |\zeta - \zeta_j|^{s_j}} \leq C \cdot \sum_{\zeta \in Z(f) \cap L_{a/2}(\zeta_i)} \frac{(1 - |\zeta|)^{p+1+\varepsilon}}{|\zeta - \zeta_i|^{s_i}} \leq C \cdot K.$$

On the other hand, if we “ignore” the product in the numerator of (1.6), we get the global bound

$$\log |f(z)| \leq \frac{K}{(1 - |z|)^p \prod_{k=1}^m |z - \xi_k|^{q_k}}, \quad z \in \mathbb{D},$$

and Theorem 0.2 in [3] gives

$$(4.7) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|)^{p+1+\varepsilon} \prod_{k=1}^m |\zeta - \xi_k|^{(q_k-1+\varepsilon)_+} \leq C \cdot K.$$

As above, the combination of (4.6) and (4.7) yields (1.7), as claimed. □

Proof of Theorem 1.3. The argument is close to the one above. Within the domain $L_a(\zeta_i)$ with

$$a := \frac{1}{2} \min_{1 \leq j \leq n} d(\zeta_j, F \cup E \setminus \{\zeta_j\}),$$

the effect of the second factor in the denominator of (1.4) is negligible. Therefore, as above in (4.6), we have with $s = \min(p, r)$

$$(4.8) \quad \sum_{\zeta \in Z(f) \cap L_{a/2}(\zeta_i)} \frac{(1 - |\zeta|)^{p+1+\varepsilon}}{d^s(\zeta, E)} \leq \sum_{\zeta \in Z(f) \cap L_{a/2}(\zeta_i)} \frac{(1 - |\zeta|)^{p+1+\varepsilon}}{|\zeta - \zeta_i|^s} \leq C \cdot K.$$

The global bound now looks as

$$(4.9) \quad \log |f(z)| \leq \frac{K}{(1 - |z|)^p d^q(z, F)}, \quad z \in \mathbb{D},$$

The Blaschke-type condition for f in (4.9) with $p = 0$ is a particular case of Theorem 3 [10]:

$$(4.10) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|) d^\rho(\zeta, F) \leq C \cdot K, \quad \rho := (q - \alpha(F) + \varepsilon)_+.$$

There is a standard way to carry the later result over to the case $p > 0$, see the proof of Theorem 0.2 in [3]. For the sake of completeness we outline the idea of this method.

Consider the sequence of functions

$$f_n(z) := f(\lambda_n z), \quad \lambda_n := 1 - 2^{-n}, \quad n \in \mathbb{N}.$$

By (4.9) and elementary inequality $d(z, F) \leq 2 d(\lambda_n z, F)$ we have

$$\log |f_n(z)| \leq \frac{2^{np+q} K}{d^q(z, F)}, \quad z \in \mathbb{D}.$$

The latter is (4.9) with $p = 0$, so, in view of (4.10),

$$(4.11) \quad \sum_{j: |\zeta_j(f)| \leq \lambda_{n-1}} (1 - |\zeta_j(f_n)|) d^\rho(\zeta_j(f_n), F) \leq C 2^{np} \cdot K,$$

where $\zeta_j(f)$, $\zeta_j(f_n)$ are the zeros of f and f_n , respectively, so $\zeta_j(f) = \lambda_n \zeta_j(f_n)$.

To obtain the lower bound of the LHS in (4.11), we note that $|\zeta_j(f)| \leq \lambda_{n-1}$ implies that

$$1 - |\zeta_j(f_n)| = 1 - \frac{|\zeta_j(f)|}{\lambda_n} \geq \frac{1 - |\zeta_j(f)|}{2}, \quad d(\zeta_j(f_n), F) \geq \frac{1}{2} d(\zeta_j(f), F),$$

and hence

$$\sum_{\lambda_n < |\zeta_j(f)| \leq \lambda_{n+1}} (1 - |\zeta_j(f)|) d^p(\zeta_j(f), F) \leq C 2^{np} \cdot K.$$

Since now

$$1 - |\zeta_j(f)| < 1 - \lambda_n = 2^{-n}, \quad (1 - |\zeta_j(f)|)^{p+\varepsilon+1} < 2^{-n(p+\varepsilon)},$$

the summation over n leads to

$$(4.12) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|)^{p+1+\varepsilon} d^{(q-\alpha(F)+\varepsilon)+}(\zeta, F) \leq C \cdot K,$$

which is the Blaschke-type condition for the functions f in (4.9) with $p > 0$. Again, a combination of (4.8) and (4.12) gives (1.8), as claimed. \square

5. Some further Blaschke-type conditions

5.1. Generalized Stolz domains

There is a seemingly more general form of the Blaschke-type condition (1.5) which states that, under assumption (1.4), for every $0 \leq \tau' < \tau < \infty$ there is a positive constant $C = C(E, F, p, q, r, \tau, \tau')$ such that

$$(5.1) \quad \sum_{\zeta \in Z(f)} (1 - |\zeta|)^{p+1+\tau} \frac{d^{(q-1+\tau)+}(\zeta, F)}{d^{\min(p,r)+\tau'}(\zeta, E)} \leq C \cdot K.$$

In fact, it is a direct consequence of (1.5) with $\varepsilon = \tau - \tau'$, since

$$\frac{1 - |\zeta|}{d(\zeta, E)} \leq 1, \quad \zeta \in \mathbb{D}.$$

However, it turns out that in some instances (5.1) holds with $\tau' = \tau$, see Corollary 5.3.

Recall the notation $\mathcal{S}_A(\zeta_0)$, $\zeta_0 \in \mathbb{T}$, $A > 0$ introduced in (2.2). In the proof of Theorem 1.4, we actually obtained a little stronger conclusion than the claimed one.

Proposition 5.1. *Let $f \in \mathcal{A}(\mathbb{D})$ be a function satisfying the assumptions of Theorem 1.4. Then for each $0 \leq \tau' < \tau < \infty$ and $\varepsilon = \tau - \tau' > 0$ there is a positive*

number $C_6 = C_6(\zeta_0, \xi_0, p, q, r, \tau, \tau')$ such that the following condition holds:

$$(5.2) \quad \sum_{\zeta \in Z(f) \cap \mathcal{S}_{1/\varepsilon}^c(\zeta_0)} \frac{(1 - |\zeta|)^{p+\tau+1} |\zeta - \xi_0|^{(q-1+\tau)_+}}{|\zeta - \zeta_0|^{\min(p,r)+\tau'}} + \sum_{\zeta \in Z(f) \cap \mathcal{S}_{1/\varepsilon}(\zeta_0)} (1 - |\zeta|)^{p+1+\varepsilon} |\zeta - \xi_0|^{(q-1+\varepsilon)_+} \leq C_6 \cdot K.$$

Obviously, inequality (5.2) reads as

$$\sum_{\zeta \in Z(f) \cap \mathcal{S}_{1/\varepsilon}^c(\zeta_0)} \frac{(1 - |\zeta|)^{p+\tau+1}}{|\zeta - \zeta_0|^{\min(p,r)+\tau'}} + \sum_{\zeta \in Z(f) \cap \mathcal{S}_{1/\varepsilon}(\zeta_0)} (1 - |\zeta|)^{p+1+\varepsilon} \leq C_6 \cdot K.$$

when $q = 0$. Of course, the above remark also holds for Theorems 1.1, 1.2.

To get sharper results we could replace summation along the Stolz angles by that along larger approach domains. For simplicity, we formulate here just the result for one point $\zeta_0 = 1$.

Theorem 5.2. *Let $f \in \mathcal{A}(\mathbb{D})$, $|f(0)| = 1$, satisfy the growth condition*

$$\log |f(z)| \leq \frac{K|1 - z|^r}{(1 - |z|)^p}, \quad z \in \mathbb{D},$$

where $0 < p < r + 1$. Then for each $\tau > 0$ there is a positive number $C_7 = C_7(p, r, \tau)$ such that

$$(5.3) \quad \sum_{\zeta \in Z(f), \frac{1-|\zeta|}{1-\zeta} > |1-\zeta|^\beta} (1 - |\zeta|) + \sum_{\zeta \in Z(f), \frac{1-|\zeta|}{1-\zeta} \leq |1-\zeta|^\beta} \frac{(1 - |\zeta|)^{p+1+\tau}}{|1 - \zeta|^{\min(p,r)+1+\tau}} \leq C_7 \cdot K,$$

where

$$\beta = \begin{cases} 1/(p + \tau), & p \leq r; \\ (r + 1 - p)/(p + \tau), & r < p < r + 1. \end{cases}$$

Corollary 5.3. *Under the same conditions,*

$$\sum_{\zeta \in Z(f)} \frac{(1 - |\zeta|)^{p+1+\tau}}{|1 - \zeta|^{\min(p,r)+\tau}} \leq C_8 \cdot K.$$

Proof. We use that $(1 - |\zeta|)/|1 - \zeta| \leq 1$ and that $1 - |\zeta| \leq 1$. Then

$$\sum_{\zeta \in Z(f), \frac{1-|\zeta|}{1-\zeta} > |1-\zeta|^\beta} \frac{(1 - |\zeta|)^{p+1+\tau}}{|1 - \zeta|^{\min(p,r)+\tau}} \leq \sum_{\zeta \in Z(f), \frac{1-|\zeta|}{1-\zeta} > |1-\zeta|^\beta} (1 - |\zeta|).$$

It remains to use (5.3). □

Proof of Theorem 5.2. Let

$$\psi(z) = \frac{1-z}{1+z}, \quad F(z) = f(\psi(z)).$$

Then F is analytic in the right half-plane \mathbb{C}_r and

$$(5.4) \quad \log |f(z)| \leq C'K \cdot \frac{|z|^r}{x^p}, \quad z \in \mathbb{C}_r, |z| < C,$$

where $z = x + iy$, $C > 1$ is arbitrary and C' depends on p, r, C .

Let $\lambda > 1$ be fixed later on. Consider the domain

$$\Omega_0 = \{x + iy : x > |y|^\lambda\}.$$

Let ϕ_0 be a conformal map of Ω_0 onto \mathbb{C}_r such that $\phi_0(\Omega_0 \cap \mathbb{R}) = \mathbb{C}_r \cap \mathbb{R}$. To obtain good asymptotic information on ϕ_0 at 0, we use the results of Warschawski [28] (see also [24], Theorem 11.16). For some C and C' depending only on λ we obtain

$$(5.5) \quad 0 < C < \frac{|\phi_0(z)|}{|z|} < C' < \infty, \quad z \in \Omega_0, |z| < 1.$$

Furthermore, the same results show that given $\gamma \in (0, 1]$ we have

$$(5.6) \quad 0 < C < \frac{\operatorname{Re} \phi_0(x + iy)}{x} < C' < \infty, \quad x > \max(\gamma|y|, 2|y|^\lambda), |x + iy| < 1,$$

with C, C' depending only on γ and λ .

Next, we need a similar estimate for

$$(5.7) \quad 2|y|^\lambda \leq x < \gamma|y|, \quad |x + iy| < \frac{1}{4}.$$

For $y \in [-1/4, 1/4]$ we consider the points

$$\begin{aligned} A &= |y|^\lambda/2 + i(y - \gamma|y|), & B &= |y|^\lambda/2 + i(y + \gamma|y|), \\ A' &= 3|y|^\lambda/2 + i(y - \gamma|y|), & B' &= 3|y|^\lambda/2 + i(y + \gamma|y|), \\ A'' &= 2|y| + i(y - \gamma|y|), & B'' &= 2|y| + i(y + \gamma|y|) \end{aligned}$$

and the rectangles $ABB''A''$, $A'B'B''A''$, see Figure 3. From now on we fix $\gamma = \gamma(\lambda)$ as the maximal number in $(0, 1]$ such that

$$A'B'B''A'' \subset ABB''A'' \cap \overline{\Omega_0}$$

for all $y \in [-1/4, 1/4]$.

Fix $x + iy$ satisfying (5.7), the corresponding points A, B, A', B', A'', B'' and the rectangles $ABB''A''$, $A'B'B''A''$. Then

$$[A'', B''] \subset \{x + iy : x > \max(\gamma|y|, 2|y|^\lambda), |x + iy| < 1\}.$$




FIGURE 3. Rectangles $ABB''A''$, $A'B'B''A''$, and the part of the boundary $\partial\Omega_0 \cap (ABB'A')$.

Set $u = \text{Re } \phi_0$. By (5.5) and (5.6) we have

$$\begin{aligned} 0 \leq u(w) &\leq C''|y|, & w \in ABB''A'' \cap \Omega_0, \\ C''|y| &\leq u(w), & w \in [A'', B''], \end{aligned}$$

is this correct?
or just C'' ?

with C'' 's depending only on λ .

Since $0 < x < |y|$,

$$\begin{aligned} d(x + iy, [A', B']) &\geq x/4, \\ d(x + iy, [A', A'']) &= d(x + iy, [B', B'']) = \gamma|y|, \\ d(x + iy, [A'', B'']) &\leq 2|y|, \end{aligned}$$

we have written these
inequalities in display
mode, OK?

an elementary estimate of harmonic measure shows that

$$\omega(x + iy, [A'', B''], A'B'B''A'') \geq C'' \cdot \frac{x}{|y|},$$

with C'' depending only on λ . Hence,

$$u(x + iy) \geq C \cdot x.$$

Since $0 < x < |y|$,

$$\begin{aligned} d(x + iy, [A, B]) &\leq x, \\ d(x + iy, [A, A'']) &= d(x + iy, [B, B'']) = \gamma|y|, \\ d(x + iy, [A'', B'']) &\geq |y|, \end{aligned}$$

another elementary estimate of harmonic measure gives that

$$\begin{aligned} \omega(x + iy, \partial(ABB''A'' \cap \Omega_0) \setminus \partial\Omega_0, ABB''A'' \cap \Omega_0) \\ \leq \omega(x + iy, \partial(ABB''A'') \setminus [A, B], ABB''A'') \leq C'' \cdot \frac{x}{|y|}, \end{aligned}$$

with C'' depending only on λ . Hence,

$$u(x + iy) \leq C \cdot x.$$

As a result, we obtain

$$(5.8) \quad 0 < C < \frac{\operatorname{Re} \phi_0(x + iy)}{x} < C' < \infty, \quad x \geq 2|y|^\lambda, |x + iy| < 1,$$

with C, C' depending only on λ .

Now, for $n \geq 1$ we define

$$\Omega_n = \{x + iy : x > 2^{-n}|y|^\lambda\},$$

and $\phi_n : \Omega_n \mapsto \mathbb{C}_r$,

$$\phi_n(z) = 2^{n/(\lambda-1)}\phi_0(2^{-n/(\lambda-1)}z).$$

By (5.5) and (5.8), for some C, C' and for $n \geq 1, z \in \Omega_n, |z| < 1$ we have

$$\begin{aligned} 0 < C < \frac{|\phi_n(z)|}{|z|} < C' < \infty, \\ 0 < C < \frac{\operatorname{Re} \phi_n(x + iy)}{x} < C' < \infty. \end{aligned}$$

Next, we define

$$G_n = F \circ \phi_n^{-1}, \quad n \geq 1.$$

Then $|G_n(\phi_n(1))| = 1$. Set $Q = \max(2|\phi_n(1)|, 1)$. By (5.4) we have

$$\begin{aligned} \log |G_n(iy)| &\leq CK \cdot 2^{np}|y|^{r-\lambda p}, \quad y \in [-Q, Q], \\ \log |G_n(e^{i\theta}Q)| &\leq CK \cdot 2^{np}, \quad \theta \in [-\pi/2, \pi/2], \end{aligned}$$

with C depending only on λ .

From now on we suppose that $r - \lambda p > -1$. By the Poisson–Jensen formula in the right half-disk $\{z \in \mathbb{C}_r : |z| < Q\}$ we obtain that

$$\sum_{G_n(x+iy)=0, |x+iy|<1/2} x \leq CK \cdot 2^{np}, \quad n \geq 1,$$

and hence,

$$\sum_{F(x+iy)=0, x>2^{1-n}|y|^\lambda, |x+iy|<C} x \leq C'K \cdot 2^{np}, \quad n \geq 1,$$

with C and C' depending only on λ, p, r . Theorem 1.4 implies that

$$\sum_{F(x+iy)=0, x>2^{1-n}|y|^\lambda, C\leq|x+iy|<1} x \leq C'K \cdot 2^{np}, \quad n \geq 1,$$

with C and C' depending only on λ, p, r . Hence,

$$\sum_{F(x+iy)=0, x>2^{1-n}|y|^\lambda, |x+iy|<1} x \leq CK \cdot 2^{np}, \quad n \geq 1,$$

with C depending only on λ, p, r .

Let $\delta > 1$. Then

$$\sum_{F(x+iy)=0, x>|y|^\lambda, |x+iy|<1} x + \sum_{F(x+iy)=0, x\leq|y|^\lambda, |x+iy|<1} \frac{x^{1+\delta p}}{|x+iy|^{\delta\lambda p}} \leq CK,$$

with C depending only on λ, p, r, δ . If $p \leq r$, then, given $\tau > 0$, we can choose $\delta = 1 + \tau/p, \lambda = 1 + 1/(p + \tau)$ to get

$$\sum_{F(x+iy)=0, x>|y|^\lambda, |x+iy|<1} x + \sum_{F(x+iy)=0, x\leq|y|^\lambda, |x+iy|<1} \frac{x^{p+1+\tau}}{|x+iy|^{p+1+\tau}} \leq CK.$$

If $r < p < r+1$, then, given $\tau > 0$, we can choose $\delta = 1 + \tau/p, \lambda = (r+1+\tau)/(p+\tau)$ to get

$$\sum_{F(x+iy)=0, x>|y|^\lambda, |x+iy|<1} x + \sum_{F(x+iy)=0, x\leq|y|^\lambda, |x+iy|<1} \frac{x^{p+1+\tau}}{|x+iy|^{r+1+\tau}} \leq CK.$$

Returning to the zeros of f and estimating those far from the point 1 as in the proof of Theorem 1.4 we obtain (5.3). □

5.2. Upper half-plane and plane with a cut

A version of Theorem 1.2 for the upper half-plane looks as follows. We use a convenient notation

$$\{u\}_{c,\varepsilon} := (u_- - 1 + \varepsilon)_+ - \min(c, u_+), \quad c \geq 0, \quad \varepsilon > 0.$$

Recall that $u_+ = \max(u, 0), u_- = \max(-u, 0)$, and so $u = u_+ - u_-$.

Theorem 5.4. *Let $X = \{x_j\}_{j=1}^n$ and $X' = \{x'_k\}_{k=1}^m$ be two disjoint finite sets of distinct points on the real line. Let $g \in \mathcal{A}(\mathbb{C}_+)$, $|g(i)| = 1$, satisfy the growth condition*

$$(5.9) \quad \log |g(w)| \leq K \frac{(1 + |w|)^{2b}}{(\operatorname{Im} w)^a} \frac{\prod_{j=1}^n |w - x_j|^{c_j}}{\prod_{k=1}^m |w - x'_k|^{d_k}}, \quad w \in \mathbb{C}_+,$$

and $a, b, c_j, d_k \geq 0$. Denote

$$l := 2a - 2b - \sum_{j=1}^n c_j + \sum_{k=1}^m d_k = l_+ - l_-.$$

Then for each $\varepsilon > 0$ there exists a positive number $C_9 = C_9(X, X', a, b, c_j, d_k, \varepsilon)$ such that the following Blaschke-type condition holds:

$$(5.10) \quad \sum_{\zeta \in Z(g)} \frac{(\operatorname{Im} \zeta)^{a+1+\varepsilon}}{(1 + |\zeta|)^{l_1}} \frac{\prod_{k=1}^m |\zeta - x'_k|^{(d_k-1+\varepsilon)_+}}{\prod_{j=1}^n |\zeta - x_j|^{\min(a, c_j)}} \leq C_9 \cdot K,$$

where the parameter l_1 is defined by the relation

$$l_1 := 2(a + 1 + \varepsilon) + \{l\}_{a, \varepsilon} - \sum_{j=1}^n \min(a, c_j) + \sum_{k=1}^m (d_k - 1 + \varepsilon)_+.$$

Proof. Since the result follows directly from Theorem 1.2, we give only a sketch of the proof. Consider the standard conformal mappings

$$(5.11) \quad z = z(w) = \frac{w - i}{w + i} : \mathbb{C}_+ \rightarrow \mathbb{D}, \quad w = w(z) = i \frac{1 + z}{1 - z} : \mathbb{D} \rightarrow \mathbb{C}_+,$$

and the following elementary relations between the corresponding quantities in \mathbb{C}_+ and \mathbb{D} :

$$\frac{2}{1 + |w|} \leq |1 - z| \leq \frac{2\sqrt{2}}{1 + |w|}, \quad \frac{2 \operatorname{Im} w}{(1 + |w|)^2} \leq 1 - |z| \leq \frac{8 \operatorname{Im} w}{(1 + |w|)^2}.$$

We have

$$|w - x_j| = \frac{2|z - \zeta_j|}{|1 - z||1 - \zeta_j|}, \quad \frac{2|w - x_j|}{(1 + |w|)|x_j + i|} \leq |z - \zeta_j| \leq \frac{2\sqrt{2}|w - x_j|}{(1 + |w|)|x_j + i|}$$

with $\zeta_j = z(x_j)$. Similar inequalities hold for $|w - x'_k|$ and $|z - z(x'_k)|$. Then, we map \mathbb{C}_+ onto \mathbb{D} using $w(z)$ defined in (5.11), and rewrite inequality (5.9) in terms of $z \in \mathbb{D}$. To complete, we apply Theorem 1.2 and go back to \mathbb{C}_+ using $z(w)$ defined in (5.11). \square

In view of applications to the spectral theory we give yet another version of Theorem 1.2 related to the domain $\mathbb{C} \setminus \mathbb{R}_+$.

Theorem 5.5. *Let $T = \{t_j\}_{j=1}^n$ and $T' = \{t'_k\}_{k=1}^m$ be two disjoint finite sets of distinct positive numbers. Let $h \in \mathcal{A}(\mathbb{C} \setminus \mathbb{R}_+)$, $|h(-1)| = 1$, satisfy the growth condition*

$$\log |h(\lambda)| \leq \frac{K}{|\lambda|^r} \frac{(1 + |\lambda|)^b}{d^a(\lambda, \mathbb{R}_+)} \frac{\prod_{j=1}^n |\lambda - t_j|^{c_j}}{\prod_{k=1}^m |\lambda - t'_k|^{d_k}}, \quad \lambda \in \mathbb{C} \setminus \mathbb{R}_+,$$

and $a, b, c_j, d_k \geq 0, r \in \mathbb{R}$. Denote

$$s := 3a - 2b + 2r - 2 \sum_{j=1}^n c_j + 2 \sum_{k=1}^m d_k = s_+ - s_-.$$

Then for each $\varepsilon > 0$ there is a positive number C which depends on all parameters involved such that the following inequality holds:

$$(5.12) \quad \sum_{\zeta \in Z(h)} d^{a+1+\varepsilon}(\zeta, \mathbb{R}_+) \frac{|\zeta|^{s_1}}{(1+|\zeta|)^{s_2}} \cdot \frac{\prod_{k=1}^m |\zeta - t'_k|^{(d_k-1+\varepsilon)_+}}{\prod_{j=1}^n |\zeta - t_j|^{\min(a, c_j)}} \leq C \cdot K,$$

where the parameter s_1, s_2 are defined by the relations

$$s_1 := \frac{\{-2r - a\}_{a, \varepsilon} - a - 1 - \varepsilon}{2},$$

$$s_2 := a + 1 + \varepsilon + \frac{\{-2r - a\}_{a, \varepsilon} + \{s\}_{a, \varepsilon}}{2} - \sum_{j=1}^n \min(a, c_j) + \sum_{k=1}^m (d_k - 1 + \varepsilon)_+.$$

The result is a direct consequence of Theorem 5.4 applied to the function $g(w) := h(w^2)$, $w \in \mathbb{C}_+$, and the elementary inequalities

$$|w| \operatorname{Im} w \leq d(w^2, \mathbb{R}_+) \leq 2|w| \operatorname{Im} w, \quad w \in \mathbb{C}_+.$$

Acknowledgments. Recently, Peter Yuditskii has celebrated his 60th anniversary. All three authors are Peter's friends, LG is his former associate at ILTPE in Kharkov, and SK is his former Ph. D. student. We would like to take this opportunity to warmly wish Peter many more years of flourishing scientific (and musical) activity as well as deep mathematical insights.

We thank D. Tulyakov for useful discussions. A part of this research was done during L. Golinskii's visit to University of Bordeaux in the framework of "IdEx Invited Scholar Programm". He gratefully acknowledges the financial support of the IdEx as well as the hospitality of the institution.

References

- [1] AHERN, P. AND CLARK, D.: On inner functions with B^p derivative. *Michigan Math. J.* **23** (1976), no. 2, 107–118.
- [2] BLASCHKE, W.: Eine Erweiterung des Satzes von Vitali über Folgen analytischer Funktionen. *S.-B. Sächs Akad. Wiss. Leipzig Math.-Natur. Kl.* **67** (1915), 194–200.
- [3] BORICHEV, A., GOLINSKII, L. AND KUPIN, S.: A Blaschke-type condition and its application to complex Jacobi matrices. *Bull. Lond. Math.Soc.* **41** (2009), no. 1, 117–123.
- [4] CUENIN, J. C., LAPTEV, A., AND TRETTER, C.: Eigenvalue estimates for non-selfadjoint Dirac operators on the real line. *Ann. Henri Poincaré* **15** (2014), no. 4, 707–736.

- [5] DEMUTH, M., HANSMANN, M. AND KATRIEL, G.: On the discrete spectrum of non-selfadjoint operators. *J. Funct. Anal.* **257** (2009), no. 9, 2742–2759.
- [6] DEMUTH, M., HANSMANN, M. AND KATRIEL, G.: Lieb–Thirring type inequalities for Schrödinger operators with a complex-valued potential. *Integral Equations Operator Theory* **75** (2013), no. 1, 1–5.
- [7] DUBUISSON, C.: On quantitative bounds on eigenvalues of a complex perturbation of a Dirac operator. *Integral Equations Operator Theory* **78** (2014), no. 2, 249–269.
- [8] DUBUISSON, C.: Notes on Lieb–Thirring type inequalities for a complex perturbation of a fractional Schrödinger operator. *Zh. Mat. Fiz. Anal. Geom.* **11** (2015), no. 3, 245–266.
- [9] FAVOROV, S. AND GOLINSKII, L.: A Blaschke-type condition for analytic and subharmonic functions and application to contraction operators. In *Linear and complex analysis*, 37–47. Amer. Math. Soc. Transl., Ser. 2, 226, Amer. Math. Soc. Providence, RI, 2009.
- [10] FAVOROV, S. AND GOLINSKII, L.: Blaschke-type conditions for analytic and subharmonic functions in the unit disk: local analogs and inverse problems. *Comput. Methods Funct. Theory* **12** (2012), no. 1, 151–166.
- [11] FRANK, R. AND SABIN, J.: Restriction theorems for orthonormal functions, Strichartz inequalities, and uniform Sobolev estimates. *Amer. J. Math.* **139** (2017), no. 6, 1649–1691.
- [12] FRANK, R.: Eigenvalue bounds for Schrödinger operators with complex potentials. III. *Trans. Amer. Math. Soc.* **370** (2018), no. 1, 219–240.
- [13] GOLINSKII, L. AND KUPIN, S.: A Blaschke-type condition for analytic functions on finitely connected domains. Applications to complex perturbations of a finite-band selfadjoint operator. *J. Math. Anal. Appl.* **389** (2012), no. 2, 705–712.
- [14] GOLINSKII, L. AND KUPIN, S.: On discrete spectrum of complex perturbations of finite band Schrödinger operators. In *Recent trends in analysis*, 113–121. Theta Ser. Adv. Math. 16, Theta, Bucharest, 2013.
- [15] GOLINSKII, L. AND KUPIN, S.: On complex perturbations of infinite band Schrödinger operators. *Methods Funct. Anal. Topology* **21** (2015), no. 3, 237–245.
- [16] GOLUBEV, V.: *Oднозначные аналитические функции. Автоморфные функции.* (Russian) [Single-valued analytic functions. Automorphic functions]. Gosudarstv. Izdat. Fiz. Mat. Lit., Moscow, 1961.
- [17] HANSMANN, M. AND KATRIEL, G.: Inequalities for the eigenvalues of non-selfadjoint Jacobi operators. *Complex Anal. Oper. Theory* **5** (2011), no. 1, 197–218.
- [18] KORENBLUM, B.: An extension of the Nevanlinna theory. *Acta Math.* **135** (1975), no. 3-4, 187–219.
- [19] KORENBLUM, B.: A Beurling-type theorem. *Acta Math.* **138** (1976), no. 3-4, 265–293.
- [20] LAPTEV, A. AND SAFRONOV, O.: Eigenvalue estimates for Schrödinger operators with complex potentials. *Comm. Math. Phys.* **292** (2009), no. 1, 29–54.
- [21] LAVRENTIEV, M. A. AND CHABAT, B. V.: *Методы теории функций комплексного переменного.* (Russian). Second edition. Nauka, Moscow, 1987. French translation: *Méthodes de la théorie des fonctions d’une variable complexe.* Editions Mir, Moscou, 1977.

[11] and [12]
updated, please
check

- [22] LEVIN, B. YA.: *Distribution of zeros of entire functions*. Translations of Mathematical Monographs 5, American Mathematical Society, Providence, 1980.
- [23] NEVANLINNA, R.: Über beschränkte analytische Funktionen. *Ann. Acad. Sci. Fenn. Ser. A* **32** (1929), no. 7.
- [24] POMMERENKE, CH.: *Boundary behaviour of conformal maps*. Grundlehren der Mathematischen Wissenschaften 299, Springer-Verlag, Berlin, 1992.
- [25] SAMBOU, D.: Lieb–Thirring type inequalities for non-self-adjoint perturbations of magnetic Schrödinger operators. *J. Funct. Anal.* **266** (2014), no. 8, 5016–5044.
- [26] SEIP, K.: An extension of the Blaschke condition. *J. London Math. Soc. (2)* **51** (1995), no. 3, 545–558.
- [27] SMIRNOV, V.: Sur les valeurs limites des fonctions, régulière à l’intérieur d’un cercle. *J. Soc. Phys. Math. Léningrade* **2** (1929) no. 2, 22–37.
- [28] WARSCHAWSKI, S. E.: On conformal mapping of infinite strips. *Trans. Amer. Math. Soc.* **51** (1942), 280–335.

Received April 4, 2016.

ALEXANDER BORICHEV: Aix Marseille Université, CNRS, Centrale Marseille, I2M, 39 rue F. Joliot-Curie, 13453 Marseille, France.
E-mail: alexander.borichev@math.cnrs.fr

check emails and
addresses

LEONID GOLINSKII: Mathematics Division, ILTPE, 47 Science ave., 61103 Kharkov, Ukraine.
E-mail: golinskii@ilt.kharkov.ua

STANISLAV KUPIN: IMB, CNRS, Université de Bordeaux, 351 ave. de la Libération, 33405 Talence Cedex, France.
E-mail: skupin@math.u-bordeaux.fr